## Enhancing psychological support and communication in intensive care unit data transmission

Giacomo Fiacco, Gennaro Catone

Department of Educational, Psychological and Communication Sciences, University Suor Orsola Benincasa, Naples, Italy

Dear Editor.

We have read with great attention and interest the article "Clinical data transmission in intensive care unit: what have we learned from COVID-19?" by Pota *et al.* We believe that the authors have addressed a very important clinical and ethical aspect regarding the transmission of clinical data. The pandemic has taught us that unforeseen emergent events are still possible and that the global healthcare and economic systems may not be adequately prepared to respond. We agree that, to date, there is no precise regulation on the transmission of patient-related clinical data to families. This includes challenges such as identifying the appropriate recipient, the lack of qualified personnel for medical communication, and the use of uncertified channels that neglect data protection regulations. Additionally, we wish to draw attention to another crucial aspect: psychological support. Healthcare personnel often find themselves communicating severe clinical evalua-

Correspondence: Giacomo Fiacco, Department of Educational, Psychological and Communication Sciences, University Suor Orsola Benincasa, Naples, Italy E-mail: giacomofcc@outlook.com

Key words: Covid-19, communication, intensive care, psychological support.

Conflicts of interest: the authors declare no conflict of interest.

Ethics approval and consent to participate: not applicable.

Informed consent: not applicable.

Patient consent for publication: not applicable.

Availability of data and materials: not applicable.

Received: 29 June 2024. Accepted: 5 July 2024.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2024 Licensee PAGEPress, Italy Advances in Anesthesia and Pain Medicine 2024; 2:44 doi:10.4081/aapm.44

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0). tions and sometimes unfavorable treatment outcomes. Some of the main challenges faced daily by operators in intensive care contexts involve continuous exposure to severe physical and psychological suffering, navigating uncertainty and unpredictability, and balancing the needs of the patient with those of their family. The complexity of current technological aspects introduces new challenges in this scenario.<sup>2</sup> The lack of in-person communication could weaken the trust and empathy that should exist between healthcare personnel and the family.

Providing adequate information requires communication skills, for which many healthcare professionals have not received proper training.3 Effective communication between healthcare personnel and family members involves creating a space for emotional expression and exchange. It is particularly important for the person delivering clinical news to have competencies in conveying clinical data. Attention to this dimension is a significant preventive factor for family health, as it has been demonstrated that post-traumatic stress, anxiety, and depression decrease following interventions to improve communication in intensive care units.4 This issue is exacerbated by non-verbal communication and the lack of adequate skills in defining the clinical situation. Additionally, the COVID-19 pandemic has had significant effects on the mental well-being of many people, regardless of the clinical severity of the infection.4 Therefore, the concept of "patient" could be extended to include family members, often also COVID-19 positive, who, despite not having severe clinical symptoms requiring hospitalization, are susceptible to anxiety and depression. This makes them even more sensitive to receiving critical news about a family member.

All these aspects highlight the necessity of involving psychologists as an integral part of clinical data transmission. Early psychological assistance allows individuals to process traumatic experiences; without processing, individuals are almost always condemned to the deleterious effects of Post-Traumatic Stress Disorder, which do not resolve naturally but tend to become chronic. Such support would be desirable in in-person meetings but is, in our opinion, essential when communication occurs through multimedia channels. The COVID-19 pandemic served as an important prototype for emergencies. Critically analyzing all processes enables the implementation of protocols that reduce issues and prevent previously encountered errors.

## References

- Pota V, Coletta F, Visconti A, et al. Clinical data transmission in intensive care unit: what have we learned from COVID-19?. Acute Care Med Surg Anesth 2023;1:26.
- 2. Stayt LC. Nurses' experiences of caring for families with rel-





## **Letter to the Editor**

- atives in intensive care units. J Adv Nurs 2007;57:623-30.
- 3. Velasco Bueno JM, La Calle GH. Humanizing intensive care: from theory to practice. Crit Care Nurs Clin North Am 2020;32: 135-47.
- 4. Lautrette A, Darmon M, Megarbane B, et al. A communication strategy and brochure for relatives of patients dying in the ICU. N Engl J Med 2007;356:469-78. Erratum in: N Engl J Med 2007;357:203.



