

Efficacy of repeated pronations in patients affected by COVID-19-related acute respiratory distress syndrome: think it twice!

Andrea Sica, Luca Gobbi, Daniele Bellantonio, Silvia Passero, Lorenzo Viola

Anesthesia and Intensive Care Unit, Bufalini Hospital, Cesena, Italy

Abstract

Prone position is widely used to ameliorate gas exchange in severe COVID-19-related ARDS (CARDS), through its beneficial effects on shunt and dead space. However, the effectiveness of repeated pronation has never been explored in this category of

patients. We then retrospectively analyzed the changes of the alveolar-arterial oxygen gradient ($A-aO_2$ grad), as shunt index, and the ratio $EtCO_2/p_aCO_2$ ratio, as dead space index, during repeated pronation cycles in 7 patients with CARDS admitted to our intensive care unit. The $A-aO_2$ grad decreased significantly more during the first pronation than the second (-43.6% vs -12.2% - $p0.008$) and, similarly, the improvement was maintained only after the first supination (-26.2% vs +9% - $p0.04$). The $EtCO_2/p_aCO_2$ ratio showed similar behavior but did not achieve statistical significance. Considering our findings, with the inherent limitations of the study, since pronation entails risks for the patient, as well as requiring a lot of effort from the nursing staff, further caution appears to be necessary in indicating pronation.

Correspondence: Andrea Sica, Anesthesia and Intensive Care Unit, Bufalini Hospital, viale Ghirelli 286, 47521 Cesena (FC), Italy.
Tel.: +39.0547.352818; Fax: +39.0547.645008.
E-mail: andrea.sica@auslromagna.it

Key words: COVID-19; pronation; pulmonary shunt; ventilation/perfusion mismatch; ARDS; CARDS.

Contributions: LV, AS and LG conceived and wrote the manuscript; DB, GO and SP reviewed manuscript and literature; all authors read and approved the final work.

Conflict of interest: the Authors declare no conflict of interest.

Ethics approval: this study was approved by the Ethical Committee of AUSL Romagna. The study is conformed with the Helsinki Declaration of 1964, as revised in 2013, concerning human and animal rights.

Informed consent and consent for publication: the data was collected retrospectively and no consent was required.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

Received: 3 March 2023.

Accepted: 24 March 2023.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Advances in Anesthesia and Pain Medicine 2023; 1:9

doi:10.4081/aapm.9

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Introduction

As part of management strategies for severe respiratory failure, prone positioning has been shown to improve the mismatch between ventilation and perfusion (V/Q) and, consequently, gas exchanges.¹ Dependent lung tissue is recruited and lung secretion drainage is promoted.²⁻⁴

Several multicenter studies and meta-analyses have shown that the application of long sessions in the prone position, together with lower tidal volume and continuous intravenous infusion of muscle relaxant, are the most important strategies leading to a reduction in overall mortality in patients with ARDS.^{5,6}

However, the mechanisms leading to these effects are not fully understood.

The advantages of prone positioning are widely exploited for the treatment of COVID-19-related acute respiratory distress syndrome;^{7,8} however, little is known about the efficacy of repeated cycles on lung shunt and dead space. Our aim is to analyze the efficacy of repeated pronation in early COVID-19-related pneumonia.

Materials and Methods

We conducted a retrospective study on patients admitted in the ICU of Bufalini Hospital in Cesena, Italy, from 03/03/2020 to 09/05/2020. We enrolled mechanically ventilated adult patients who underwent at least two cycles of prono-supination within the first five days of ICU hospitalization. As standard of care, we set minimum F_iO_2 to achieve $p_aO_2 \geq 70$ mmHg and minute ventilation to maintain $pH \geq 7.25$ or $p_aCO_2 < 70$ mmHg. We registered end-tidal CO_2 ($EtCO_2$) and arterial blood gas analysis just before every decubitus change and one hour after supination. Finally, we calculated shunt in terms of alveolar-arterial oxygen gradient ($A-aO_2$ grad), computed as

$$[(F_{iO_2}) \times (\text{atmospheric pressure} - H_2O \text{ pressure}) - (p_aCO_2/0.8)] - p_aO_2$$

EtCO₂/p_aCO₂ ratio (EtCO₂/p_aCO₂R) was used as a surrogate of pulmonary dead space.

Results

Data were fully available for seven patients; only two of them underwent more than two pronation-supination cycles, so these records were excluded from the analysis. Median pronation time was 23 (IQR 6) hours for the first cycle and 18 (IQR 4.5) hours for the second one (p=0.13). In order to better understand the effect of the single pronation maneuver, we have analyzed the variation of the indices cited above during the pronation (difference between end of supination and end of pronation) and between the same (difference between end of supination and one hour after end of pronation). Along the first pronation, median A-aO₂grad decreased from 369.3 (IQR 293) to 208.4 (IQR 146.4) mmHg and EtCO₂/p_aCO₂R increased from 0.72 (IQR 0.26) to 0.84 (IQR 0.08); one hour after supination A-aO₂grad was 272.6 mmHg and EtCO₂/p_aCO₂R was 0.83. The second cycle led to a median A-aO₂grad reduction from 298.8 (IQR 95.2) to 262.4 (IQR 147.6) mmHg, while EtCO₂/p_aCO₂R raised from 0.81 (IQR 0.06) to 0.84 (IQR 0.11); A-aO₂grad and EtCO₂/p_aCO₂R were, respectively, 325.8 mmHg and 0.76 one hour after supination. The major procedure-related complications were pneumothorax (1 case) and accidental ventilator disconnection with sudden severe desaturation (1 case), the former drained and the latter resolved with a recruitment maneuver.

Discussion

The inflammatory state consequent to SARS-CoV-2 infection leads to a pulmonary epithelial and endothelial damage, with deleterious effects on gas exchange and pulmonary vascular resistance (*i.e.*, V/Q ratio).⁹ Pulmonary shunt and respiratory dead space are difficult to evaluate bedside, as their quantification requires advanced instruments and highly invasive devices. Hence, we focused on the variation of EtCO₂/p_aCO₂R for respiratory dead space and A-aO₂grad for pulmonary shunt, which are more clinically suitable.^{10,11}

A-aO₂grad decreased by 43.6% at the end of the first pronation, while the reduction was 12.2% with the second cycle (p=0.008, Mann-Whitney test). This trend is confirmed by the improvement of EtCO₂/p_aCO₂R (+15.6% after the first cycle, +3.7% after the second one – p=0.16). It is interesting to note that the most marked improvement that occurs during the first pronation is reflected also at the moment of supination: the improvement of A-aO₂grad, meant as the difference between before the pronation and an hour after the supination, is significantly higher after the first cycle than the second (p=0.04) (Figure 1). Also from this point of view EtCO₂/p_aCO₂R confirms the trend improvement, although not reaching the significance (p=0.10) (Figure 2).

The lesser duration of the second pronation does not seem to be correlated to its minor efficacy; indeed, there is no correlation between the pronation length and A-aO₂grad and EtCO₂/p_aCO₂R in the first cycle (respectively, Spearman's rho 0.41 and 0.45), even less in the second one (respectively, Spearman's rho -0.19 and -0.45).




Figure 1. A-aO₂ gradient variations during the pronation and one hour after supination in the first and in the second pronation-supination maneuver.




Figure 2. $\text{EtCO}_2/\text{paCO}_2$ ratio variations during the pronation and one hour after supination in the first and in the second prono-supination maneuver.

Conclusions

The little amount of data limits the statistical significance of our study; nevertheless, our work gives new insights on the suitability of repeated pronation maneuvers. Although simple, the pronation is not free of complications and requires an additional burden in an over-ceiling work environment, such as during the COVID-19 pandemic.

References

- Gattinoni L, Busana M, Giosa L, et al. Prone positioning in Acute Respiratory Distress Syndrome. *Semin Respir Crit Care Med* 2019;40:94-100.
- Abroug F, Ouannes-Besbes L, Elatrous S, et al. The effect of prone positioning in acute respiratory distress syndrome or acute lung injury: a meta-analysis: areas of uncertainty and recommendations for research. *Intensive Care Med* 2008;34:1002-11.
- Sud S, Friedrich JO, Adhikari NK, et al. Effect of prone positioning during mechanical ventilation on mortality among patients with acute respiratory distress syndrome: a systematic review and meta-analysis. *CMAJ* 2014;186:E381-E390.
- Guérin C, Reigner J, Richard JC, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368:2159-68.
- Petrucchi N, De Feo C. Lung protective ventilation strategy for the acute respiratory distress syndrome. *Cochrane Database Syst Rev* 2013;2:CD003844.
- Brochard L, Slutsky A, Pesenti A. Mechanical ventilation to minimize progression of lung injury in acute respiratory failure. *Am J Respir Crit Care Med* 2017;195:438-442.
- Qadri SK, Ng P, Toh TSW, et al. Critically ill patients with COVID-19: a narrative review on prone position. *Pulm Ther* 2020;6:233-46.
- Scaramuzza G, Gamberini L, Tonetti T, et al. Sustained oxygenation improvement after first prone positioning is associated with liberation from mechanical ventilation and mortality in critically ill COVID-19 patients: a cohort study. *Ann Intensive Care* 2021;11:63.
- Bussani R, Schneider E, Zentilin L. Persistence of viral RNA, pneumocyte syncytia and thrombosis are hallmarks of advanced COVID-19 pathology. *E Bio Medicine* 2020;61:103-4.
- Gattinoni L, Chiumello D, Rossi S. COVID-19 pneumonia: ARDS or not? *Crit Care* 2020;24:154.
- Harris DE, Massie M. Role of alveolar-arterial gradient in partial pressure of oxygen and $\text{PaO}_2/\text{fraction of inspired oxygen}$ ratio measurements in assessment of pulmonary dysfunction. *AANA J* 2019;87:214-21.